

Quasi-normal modes of D3-brane Black Holes

Yasunari Kurita*

Graduate School of Human and Environmental Studies, Kyoto University, Kyoto 606-8501, Japan

Masa-aki Sakagami†

Department of Fundamental Sciences, FIHS, Kyoto University, Kyoto, 606-8501, Japan

(Dated: February 1, 2008)

We investigate a method to evaluate quasi-normal modes of D3-brane black holes by wave interpretation of fields on D3-brane based on the Feynman's space-time approach. We perturbatively solve the wave equation which describes propagation of a dilaton wave in a bulk space and its interaction with the D3-brane. The obtained condition for the quasi-normal modes are qualitatively equivalent to that evaluated in the usual scattering of the dilaton in the black 3-brane spacetime in the corresponding supergravity description.

PACS numbers: 04.70.-s, 04.50.+h, 11.25.Hf, 11.15.-q

I. INTRODUCTION

It is well known that black holes have thermodynamical properties, i.e. entropy[1, 2] and temperature[3], which can be understood in the framework of general relativity[4] and quantum theory of matter in curved spacetime[5]. In these formalisms, the entropy for a black hole is given by a quarter of its horizon area and Hawking radiation can be explained as a particle creation caused by existence of the horizon. Some years ago, in the string theory, which seems to be most promising candidate for a quantum theory of gravity, it was found that the D-brane can also describe black hole spacetime and its thermodynamical properties[6]. In this prescription, the entropy for a black hole is derived by counting the number of microscopic states on the D-brane[7], and Hawking radiation can be recognized as emission process of closed strings from the D-brane[8, 9].

As for the scattering processes of a particle or a wave in a black hole spacetime, we can observe a good agreement between a D-brane picture and an analysis based on the corresponding supergravity description. Especially the absorption cross section for a dilaton by the D3-brane in a low-energy region, which is evaluated by means of the world volume approach, has been shown to coincide with the result obtained by solving the wave equation for the dilaton field propagating in the 3-brane background[10, 11, 12]. Taking account of the above results in the scattering processes, it seems to be quite natural to pose a question whether the agreement is hold even for quasi-normal modes (QNMs) of a black hole or not.

QNMs characterize the emission of a gravitational wave which represents a response to a perturbation affected to a black hole spacetime[13, 14, 15]. As briefly reviewed in appendix A, it is obtained by solving the wave

equation on the background with the suitable boundary conditions that the flux at the horizon is ingoing and outgoing at spatial infinity. QNM is characterized by a complex frequency whose imaginary part represents the time scale in which the perturbation to the black hole spacetime decays. Since black holes are interpreted as thermal objects that are characterized by thermal quantities such as the temperature and entropy, the imaginary part of the QNM frequency can be recognized as the relaxation time within which black holes approach to the thermal equilibrium.

As was well known, the relationship between D-branes and black holes in string theory is an important precursor to the AdS/CFT correspondence. And connection between QNMs and the decay of perturbations in the dual CFT was first suggested in the work of [16] based on the numerical computation of QNMs for AdS-Schwarzschild black holes in several dimensions. QNMs of AdS black holes and BTZ black holes are investigated in [17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27] and those of near extremal black branes are found in [28]. Furthermore in [29], it was shown that the frequencies of QNMs for BTZ black holes are in exact agreement with the location of the poles of the retarded correlation function describing the linear response on the CFT side.

In the present paper, we consider the D3-brane as another example of the correspondence in QNMs. Our analyses for the D3-branes will be performed in the two different parameter regions in type IIB string theory, one is large number of the D3-branes and the other is only one D3-brane. In the first region, the classical supergravity is effective so that gravity is described by the curvature of space-time. In the later, the perturbative field theory on the D3-brane, which is embedded in a flat spacetime, is effective. In such flat spacetime, how QNMs can be calculated? This is our main interest and motivation in this paper. We will apply Feynman's space-time approach to this problem, which gives alternative intuitive methods to quantum field theories [30, 31, 32, 33]. We perturbatively solve the equation derived from the low energy effective action for the D3-brane, i.e. the Dirac-Born-

*Electronic address: kurita@phys.h.kyoto-u.ac.jp

†Electronic address: sakagami@phys.h.kyoto-u.ac.jp

Infeld action, which describes propagation of a dilaton wave in the bulk flat spacetime and its interaction with D-branes. And we apply a general condition for QNMs which is obtained in appendix A to the scattering problem of this flat D3-brane case. It is shown that the obtained condition of QNMs is qualitatively equivalent to that evaluated in the framework of the usual scattering of the dilaton wave based of the supergravity description.

A brief outline of this paper is following. In the next section, we obtain the QNMs for the 3-brane by solving the wave equation propagating on its background in classical supergravity analysis. In section 3, only one D3-brane are considered and the condition of the QNMs are obtained by the Feynman's space-time approach. These two results are compared and discussed in section 4. In appendix A, we give a brief review of QNMs. Absorption of a scalar for D3-brane is also investigated by space-time approach and the result is compared with the result of [11] in appendix C.

II. QUASI-NORMAL MODES IN SUPERGRAVITY

In this section, we evaluate the QNM of 3-brane solution of type IIB supergravity. The main analysis of this section is based on the results obtained in [12], which studies absorption probability of a dilaton by the D3-brane. We consider the dilaton as a minimally coupled scalar, which obeys the wave equation

$$\frac{1}{\sqrt{-g}}\partial_A\sqrt{-g}g^{AB}\partial_B\phi=0, \quad (A, B=0, 1, \dots, 9) \quad (1)$$

in the spacetime whose metric is given by

$$ds^2 = \left(1 + \frac{R^4}{r^4}\right)^{-1/2} (-dt^2 + dx_i dx^i) + \left(1 + \frac{R^4}{r^4}\right)^{1/2} (dr^2 + r^2 d\Omega_5^2). \quad (2)$$

The characteristic length of the 3-brane R is related to 10-dimensional gravitational coupling constant κ_{10} as follows,

$$R^4 = 4\pi g N \alpha'^2 = \frac{N \kappa_{10}}{2\pi^{5/2}}. \quad (3)$$

In the case of the scattering in the low energy region $\omega R \ll 1$, the dominant contribution to the cross section comes from spherical symmetric process so that we concentrate the radial equation for s-wave of energy ω derived from Eq.(1)

$$\left[\frac{\partial^2}{\partial r^2} + \frac{5}{r}\frac{\partial}{\partial r} + \omega^2\left(1 + \frac{R^4}{r^4}\right)\right]\phi(r) = 0. \quad (4)$$

If one performs the change of variables, $r = Re^{-z}$, $\phi(r) = e^{2z}\psi(z)$, then Eq.(4) becomes Mathieu equation,

$$\left[\frac{\partial^2}{\partial z^2} + 2(\omega R)^2 \cosh 2z - 4\right]\psi(z) = 0. \quad (5)$$

As was shown in [12], the exact solution which is ingoing at horizon ($r \rightarrow 0$) can be expressed as expansions in terms of Bessel and H ankel functions as follows,

$$\psi(z) = \sum_{n=-\infty}^{\infty} \frac{c(n+\mu)}{c(\mu)} J_n(\omega R e^{-z}) H_{n+2\mu}^{(1)}(\omega R e^z), \quad (6)$$

where the coefficients is given by

$$\begin{aligned} c(\mu) &= \frac{(\frac{\omega R}{2})^{2\mu}}{\Gamma(\mu+2)\Gamma(\mu)} v(\mu) \\ v(\mu) &= \sum_{n=0}^{\infty} (-1)^n \left(\frac{\omega R}{2}\right)^{4n} A_{\mu}^{(n)} \\ A_{\mu}^{(0)} &= 1 \\ A_{\mu}^{(q)} &= \sum_{p_1=0}^{\infty} \sum_{p_2=2}^{\infty} \cdots \sum_{p_q=2}^{\infty} a_{\mu+p_1} a_{\mu+p_1+p_2} \cdots a_{\mu+p_1+\dots+p_q} \\ a_{\mu} &= \frac{1}{\mu(\mu+1)(\mu+2)(\mu+3)}. \end{aligned} \quad (7)$$

The value of μ is determined in terms of a prescription in the standard Floquet analysis which implies

$$\frac{c(\mu)}{c(\mu-1)} \frac{c(-\mu+1)}{c(-\mu)} = 1. \quad (8)$$

The explicit expression for the first few terms of μ and $A_{\mu}^{(q)}$ is given in Appendix B. Let me introduce new variables for convenience,

$$\eta = e^{2i\pi\mu}, \quad \chi = \frac{c(-\mu)}{c(\mu)}. \quad (9)$$

With these variables, asymptotic forms of the solution (6) near the horizon ($\text{Re } z \rightarrow \infty$) is given by

$$\begin{aligned} &\sqrt{\eta} \left(\eta - \frac{1}{\eta}\right) \psi(z) \\ &\rightarrow \left(\eta - \frac{1}{\eta}\right) \sqrt{\frac{2}{\pi\omega R e^z}} e^{i(\omega R e^z - \frac{\pi}{4})}. \end{aligned} \quad (10)$$

Similarly we can obtain the asymptotic form for the spatial infinity ($\text{Re } z \rightarrow -\infty$)

$$\begin{aligned} &\sqrt{\eta} \left(\eta - \frac{1}{\eta}\right) \psi(z) \\ &\rightarrow \left(\chi - \frac{1}{\chi}\right) \sqrt{\frac{2}{\pi\omega R e^{-z}}} e^{i(\omega R e^{-z} - \frac{\pi}{4})} \\ &\quad + \left(\eta\chi - \frac{1}{\eta\chi}\right) \sqrt{\frac{2}{\pi\omega R e^{-z}}} e^{-i(\omega R e^{-z} - \frac{\pi}{4})}. \end{aligned} \quad (11)$$

From these asymptotic behavior given above, we can read off the amplitudes

$$\mathcal{A} = \left(\eta - \frac{1}{\eta}\right), \quad \mathcal{I} = \left(\eta\chi - \frac{1}{\eta\chi}\right), \quad \mathcal{R} = \left(\chi - \frac{1}{\chi}\right),$$

for the transmitted, incident and reflected waves, respectively. As we describe in Appendix A, QNMs are given by the condition,

$$\mathcal{I}/\mathcal{R} = 0. \quad (12)$$

In order to compare the obtained results (12) with the

evaluation of QNMs in the next section based on Dirac-Born-Infeld action which is the low energy effective theory of the D3-branes, we consider the QNM condition (12) in the low energy region, i.e., $\omega R \ll 1$, that can be expanded with respect to ωR as

$$\begin{aligned} \frac{\mathcal{I}}{\mathcal{R}} e^{i\beta} &= 1 + \frac{\pi^2}{512} (\omega R)^8 \left(1 + \frac{4i}{\pi} \log \frac{\omega \bar{R}}{2} \right) + \frac{\pi^2}{512 \cdot 12} (\omega R)^{12} \left(-2 \log \frac{\omega \bar{R}}{2} + \frac{4i}{\pi} (\log \frac{\omega \bar{R}}{2})^2 \right) \\ &\quad + \frac{\pi^2}{512 \cdot 72} (\omega R)^{12} \left(1 + \frac{4i}{\pi} \log \frac{\omega \bar{R}}{2} \right) + \mathcal{O}((\omega R)^{16} (\log \omega R)^3) \\ &= 0, \end{aligned} \quad (13)$$

where $\bar{R}$ is written by the characteristic length of the 3-brane R and the Euler constant γ as $\bar{R} = e^\gamma R$ and β is a phase factor,

$$\beta = \frac{2}{3} \pi \left(\frac{\omega R}{2} \right)^4 + \frac{259}{216} \pi \left(\frac{\omega R}{2} \right)^8 - \frac{22}{81} \pi^3 \left(\frac{\omega R}{2} \right)^{12} + \dots \quad (14)$$

Let us solve approximately the above QNM condition up to the lowest order $(\omega R)^8$,

$$1 + \frac{\pi^2}{512} (\omega R)^8 \left(1 + \frac{4i}{\pi} \log \frac{\omega \bar{R}}{2} \right) = 0. \quad (15)$$

In the usual evaluation to QNMs [19, 20, 21, 22, 23, 24, 25, 27], their frequencies are characterized by the curvature scale of the black hole space time, i.e. $|\omega R| \sim 1$. Contrastingly, our calculation, which is valid in the range to $|\omega R| \ll 1$, gives QNMs in the low energy region. From the observational point of view, the frequencies for QNMs obtained in this paper might not be important. However recall that our main interest is to confirm the equivalence between two different pictures, i.e. the D3-branes and black 3-branes, in the case of QNMs.

From the condition (15) in the region $|\omega R| \ll 1$, we note that its solutions lie on n -th Riemannian sheets with $n \gg 1$ in the complex ωR plane. Inserting a polar coordinate representation for $(\omega R)^8$ to Eq.(15),

$$(\omega R)^8 = r e^{i\theta}, \quad (16)$$

we obtain two real equations,

$$\begin{aligned} 0 = & 1 + \frac{\pi^2}{29} r \cos \theta \\ & - \frac{\pi r}{2^{10}} \left[(\log r + 8 \log \frac{e^\gamma}{2}) \sin \theta + \theta \cos \theta \right]. \end{aligned} \quad (17)$$

$$0 = 2\pi \sin \theta + \left[(\log r + 8 \log \frac{e^\gamma}{2}) \cos \theta - \theta \sin \theta \right]. \quad (18)$$

From Eq.(18), r is expressed in terms of θ as,

$$r = 2^8 e^{-8\gamma} \exp[(\theta - 2\pi) \tan \theta]. \quad (19)$$

So, we get the equation for θ from Eq.(17) and Eq.(19),

$$1 - \frac{\pi}{4e^{8\gamma}} e^{(\theta - 2\pi) \tan \theta} \frac{\theta - 2\pi}{\cos(\theta - 2\pi)} = 0. \quad (20)$$

Since, r must satisfy $r \ll 1$, we can see from Eq.(17) that θ must satisfy $|\theta| \gg 1$. There are two cases for the solution of Eq.(19) and Eq.(20) as follows,

$$\begin{aligned} (i) \quad & \theta > 2\pi, \quad \tan \theta < 0, \quad \cos(\theta - 2\pi) > 0 \\ \Leftrightarrow \quad & -\frac{\pi}{2} + 2n\pi < \theta < 2n\pi, \quad n \geq 2 \\ (ii) \quad & \theta < 2\pi, \quad \tan \theta > 0, \quad \cos(\theta - 2\pi) < 0 \\ \Leftrightarrow \quad & -(2n - 1)\pi < \theta < -(2n - 1)\pi + \frac{\pi}{2}, \quad n \geq 0. \end{aligned} \quad (21)$$

where n is an integer. The condition $|\theta| \gg 1$ is satisfied when $n \gg 1$. Let us consider the case (i). We put $\theta = 2n\pi - \Delta_n$ and $0 < \Delta_n < \pi/2$. Then Eq.(20) can be written as follows,

$$1 - \frac{\pi}{4e^{8\gamma}} e^{-2n\pi \tan \Delta_n} \frac{2n\pi}{\cos \Delta_n} \simeq 0. \quad (23)$$

Δ_n must be much less than unity in order that the second term of the above equation cancels the first term. It follows that,

$$1 - \frac{\pi}{4e^{8\gamma}} e^{-2n\pi \Delta_n} 2n\pi \simeq 0, \quad (24)$$

$$\Rightarrow \Delta_n \simeq \frac{1}{2n\pi} (\log n + \log \frac{\pi^2}{2e^{8\gamma}}). \quad (25)$$

In the case (ii), we put $\theta = -(2n + 1)\pi + \Delta_n$. After a similar analysis, we have the same result Eq.(25) for

Δ_n . This Δ_n are very small number as long as $n \gg 1$, so that the solution of Eq.(15) is just below the real axis of complex $(\omega R)^8$ plane. The radial parameter r can be written as,

$$r = \frac{2^9}{\pi^2 n} + \mathcal{O}\left(\frac{\log n}{n^2}\right) \quad (26)$$

The QNMs frequency are R^{-1} times eighth roots of $re^{i\theta}$.

III. THE WAVE SCATTERING OF D3-BRANE IN SPACE-TIME APPROACH

In this section, we consider the case of a small number of the D3-brane in string theory, so that the description of the system based on the field theory with Dirac-Born-Infeld action is effective one rather than that in terms of the supergravity which has been used in the previous section. For simplicity, we treat one D3-brane in flat space-time and field theory with DBI action on it. In the case of the dilaton absorption by D3-brane[10, 11, 12], the evaluation of the cross section is based on the world-volume interpretation, in which bulk dilaton field is recognized as a source of the world-volume fluctuation. In other words, it is assumed that quantum fluctuation of the bulk fields decouples and the dynamics is strictly on the brane.

As explained in the Appendix A, QNMs are determined by the condition (A10) which states no incident wave in the scattering of the dilaton field in the black hole spacetime, and the reflective amplitude is needed to evaluate QNMs frequency. The world-volume approach [10, 11, 12] do not give reflective amplitude which is

concerned to bulk propagation. Thus the world-volume approach[10, 11, 12] is not suitable for investigation of QNMs. Since the above definition of QNMs (A10) based on the scattering processes of the dilaton, we have to take account of its bulk propagation at least. In order to keep close similarity to the evaluation of QNMs in previous section based on the classical supergravity, we throw a dilaton into a D3-brane from spatial infinity, which is treated as a wave in field theory such as R.P.Feynman's space-time approach [30, 31, 32, 33]. Incident dilaton propagates in the bulk space and eventually interacts with a D3-brane. By perturbatively solving the wave equation for the dilaton, we can obtain its reflective amplitude. Another reason for using this space-time approach is that we bear its waveform in mind, which is one of the other important aspects of QNMs. The calculation based on CFT [29] might be enough to evaluate the frequencies of QNMs which are given by positions of poles of the retarded CFT correlator. However our approach can tell us actual wave forms observed at infinity. Thus it is possible to discuss relation between two pictures of black hole spacetime, i.e. the supergravity and the D-brane description, from the viewpoint not only of the frequencies of QNMs but also their wave forms. In the appendix C, for a proof of the validity of our calculation based on the Feynman's spacetime approach, the dilaton absorption cross section is evaluated by the formalism which developed in the present paper. And we show that our result agrees in the lowest order with that obtained by the world volume approach[10, 11, 12].

For a low-energy dilaton, DBI action can be expanded in Einstein frame as follows[11],

$$\begin{aligned} S_{\text{DBI}} &= -T_3 \int d^4x \sqrt{-\det \left(G_{\mu\nu}^E + \frac{e^{-\frac{\phi}{2}}}{\sqrt{T_3}} F_{\mu\nu} \right)} \\ &= \int d^4x \left(\frac{1}{4} F^2 - \frac{\sqrt{2}}{4} \kappa_{10} \phi F^2 + \frac{1}{8T_3} (F^4 - \frac{1}{2} (F^2)^2) \right) + \dots, \end{aligned} \quad (27)$$

where $F^n = F_{\mu_2}^{\mu_1} \cdots F_{\mu_n}^{\mu_1}$ and T_3 is the tension of the D3-brane. The term $F_{\mu\nu}$ is field strength of the gauge field on the 3-brane describing its longitudinal dynamics, while its transverse oscillations do not couple with dilaton. For

$N = 1$ case, gauge fields on the D3-brane are photon, and we take Coulomb gauge for it on the D3-brane. Then the action for the dilaton in the bulk and the gauge fields on the brane is given by

$$S_{\text{eff}} = -\frac{1}{2} \int d^{10}x \partial_A \phi \partial^A \phi - \frac{1}{2} \int d^{10}x \delta^{(6)}(x) \left[(1 - \sqrt{2} \kappa_{10} \phi) \partial_\mu A_i \partial^\mu A^i + \frac{1}{2T_3} (\epsilon^{ijk} \dot{A}_i \partial_j A_k) (\epsilon^{lmn} \dot{A}_l \partial_m A_n) \right], \quad (28)$$

where $i = 1, 2, 3$ and $\dot{A} = \partial_0 A$. For a low energy dilaton,

the scattering process is dominated by s-wave so that we

can put its form as $\phi = \phi(t, r)$ where $r^2 = x_4^2 + \dots + x_9^2$. The evolution of the dilaton in the bulk and the gauge field on the brane are expressed by the wave equations,

$$(-\partial_t^2 + \partial_r^2 - \frac{15}{4r^2})\varphi = -\delta(r)\frac{\kappa_{10}}{\sqrt{2\pi^3}r^{5/2}}\partial_\mu A_i \partial^\mu A^i, \quad (29)$$

$$\begin{aligned} \partial_\mu \partial^\mu A^i &= \sqrt{2}\kappa_{10}\partial_\mu(\phi\partial^\mu A^i) \\ &+ \frac{1}{T_3}\partial_0 \left[(\epsilon^{lmn}\dot{A}_l\partial_m A_n)\epsilon^{ijk}\partial_j A_k \right] \\ &+ \frac{1}{T_3}\partial_a \left[(\epsilon^{lmn}\dot{A}_l\partial_m A_n)\epsilon^{bai}\dot{A}_b \right], \end{aligned} \quad (30)$$

where we change the variable as $\varphi = r^{5/2}\phi$. We can solve these wave equations by means of perturbation with respect to interaction with gauge fields, i.e., the coupling constant κ_{10} , and four point interaction of gauge fields, whose coupling constant is $\frac{1}{T_3}$. At first, these waves are formally expanded in terms of the two coupling constants,

$$\begin{aligned} \varphi &= \varphi_{(0,0)} + \kappa_{10}\varphi_{(1,0)} + \frac{\kappa_{10}}{T_3}\varphi_{(1,1)} \\ &+ \kappa_{10}^2\varphi_{(2,0)} + \frac{\kappa_{10}^2}{T_3}\varphi_{(2,1)} + \dots, \end{aligned} \quad (31)$$

$$A_i = A_{(0,0)}^i + \kappa_{10}A_{(1,0)}^i + \frac{1}{T_3}A_{(0,1)} + \frac{\kappa_{10}}{T_3}A_{(1,1)}^i + \dots \quad (32)$$

We are interested only in response of the D3-brane to the incident dilaton wave. So we omit terms $A_{(0,1)}$ and $\phi_{(1,0)}$ in the above expansions, since these terms represent excitation which exists before arrival of the injected dilaton at 3-brane.

At (0,0)-order, we obtain wave equations

$$(-\partial_t^2 + \partial_r^2 - \frac{15}{4r^2})\varphi_{(0,0)} = 0, \quad (33)$$

$$\partial_\mu \partial^\mu A_{(0,0)}^i = 0. \quad (34)$$

which express that the dilaton wave does not interact with the D3-brane at the origin $r = 0$, and pass through it. So, regularity of the dilaton at the origin requires that the wave is reflected with the same amplitude as that of incident one. The solutions in the lowest order are given by

$$\varphi_{(0,0)} = Ce^{-i\omega t}\sqrt{r}\left[H_2^{(1)}(\omega r) + H_2^{(2)}(\omega r)\right], \quad (35)$$

$$\begin{aligned} A_{(0,0)}^i &= \int \frac{d^3k}{(2\pi)^3} \sqrt{\frac{V_3}{2k_0}} (\epsilon_{\mathbf{k}}^\alpha)^i e^{i\mathbf{k}\cdot\mathbf{x}} \\ &\times \left[e^{ik_0 t} \sum_{\alpha=1}^2 C_\alpha^* + e^{-ik_0 t} \sum_{\alpha=1}^2 C_\alpha \right]. \end{aligned} \quad (36)$$

where, $k_0 = |\mathbf{k}|$, $(\epsilon_{\mathbf{k}}^\alpha)^i$ is polarization vector, and the complex number C_α , which satisfies $|C_\alpha|^2 = 1$, represents a degree of freedom for the phase of the initial configuration of the gauge field. We prepare the wave function in

the lowest order, $A_{(0,0)}^i$, to express its vacuum fluctuation of the photon so that its normalization is taken to be as same as that in the corresponding quantum field theory.

At (1,0)-order, the wave equations for the gauge field is

$$\partial_\mu \partial^\mu A_{(1,0)}^i = -\sqrt{2}\dot{\varphi}_{(0,0)}\dot{A}_{(0,0)}^i. \quad (37)$$

Here we choose a negative energy photon $A_{(0,0)}$ in Eq.(37). Thus this equation shows that the positive energy photon $A_{(1,0)}^i$ is created from the positive energy dilaton $\varphi_{(0,0)}$ and the negative energy photon $A_{(0,0)}^i$. Because, in standard prescription of quantum field theories, a negative energy wave function is interpreted as an anti-particle which moves opposite direction, Eq.(37) describes a dilaton annihilation on the D3-brane and pair creation of photons as the calculation based on CFT in [10]. In order to construct the solution of Eq.(37), we use the retarded Green's function for 4-dimensional d'Alembertian,

$$G_A(t, \mathbf{x}; t', \mathbf{x}') = -\frac{\delta(t - t' - |\mathbf{x} - \mathbf{x}'|)}{4\pi|\mathbf{x} - \mathbf{x}'|}, \quad (38)$$

satisfying

$$\partial_\mu \partial^\mu G_A(x; x') = \delta^{(4)}(x - x'), \quad (39)$$

which leads to created photon on the D3-brane,

$$\begin{aligned} A_{(1,0)}^i &= \frac{\sqrt{2}C\omega^3}{8} \int \frac{d^3k}{(2\pi)^3} \sqrt{\frac{V_3}{2k_0}} e^{-i(\omega - k_0)t} e^{ik_i x^i} \\ &\times \left(\frac{1}{\omega} + \frac{1}{2k_0 - \omega - i\epsilon} \right) \sum_{\alpha=1}^2 C_\alpha^* (\epsilon_{\mathbf{k}}^\alpha)^i. \end{aligned} \quad (40)$$

The wave equation for the dilaton at (2,0)-order is given by

$$(-\partial_t^2 + \partial_r^2 - \frac{15}{4r^2})\varphi_{(2,0)} = -\frac{2\delta(r)}{\sqrt{2\pi^3}r^{5/2}}\partial_\mu A_{(0,0)}^i \partial^\mu A_{(1,0)}^i. \quad (41)$$

The homogeneous part of the solution for the above equation should be zero, since there is no incident dilaton at (2,0)-order. And the Eq.(41) represents the process that the excited gauge field $A_{(1,0)}^i$ and positive energy photon $A_{(0,0)}^i$ annihilate to emit a dilaton wave away from the D3-brane. We can construct retarded Green's function for dilaton,

$$\begin{aligned} G_R(t, r; t', r') &= \frac{\pi}{2i}\theta(t - t') \int \frac{d\omega}{2\pi} e^{-i\omega(t-t')} \\ &\times \left[\theta(r - r')H_2^{(1)}(\omega r)J_2(\omega r')\sqrt{rr'} + (r \leftrightarrow r') \right], \end{aligned} \quad (42)$$

which satisfies the equation,

$$(-\partial_t^2 + \partial_r^2 - \frac{15}{4r^2})G_R(t, r; t', r') = \delta(t - t')\delta(r - r'). \quad (43)$$




FIG. 1: The 1-loop self energy for the dilaton. A straight and wavy lines represent propagating dilaton and photon, respectively

The explicit expression (43) shows that this Green's function is regular at the origin $r = 0$ and obeys the outgoing condition at the spatial infinity $r \rightarrow \infty$. Then the $(2, 0)$ dilaton which is integrated over the D3-brane is given by

$$\begin{aligned}\bar{\varphi}_{(2,0)} &= \frac{1}{V_3} \int_{V_3} d^3x \varphi_{(2,0)} \\ &= -\frac{C \omega^8}{32 \cdot 64\pi^3} e^{-i\omega t} H_2^{(1)}(\omega r) \sqrt{r} \\ &\quad \times \left(1 + \frac{i}{\pi} \log \frac{\omega}{\Lambda}\right),\end{aligned}\quad (44)$$

where Λ is a cut-off factor. The Eq.(41) tells us that $\bar{\varphi}_{(2,0)}$ is created through annihilation of two photons, $A_{(0,0)}^i$ and $A_{(1,0)}^i$, which are described by Eq.(37). As discussed before, it can be interpreted that two photons are emerge via the pair creation process. Thus in the framework of quantum field theories, the external line corrected by the 1-loop self energy Feynman diagram depicted in Fig. 1 would correspond to the dilaton wave function $\bar{\varphi}_{(2,0)}$. We note that the field $A_{(0,0)}$ appeared in Eq.(37) and that in Eq.(41) are identical ones as shown in Fig.1, although we interpret these two photons as particle-antiparticle pair. Thus, in the evaluation of Eq.(44), momentum integrations are doubly counted so that we have introduced a symmetric factor $1/2$. The divergent contribution to the $(2, 0)$ -order dilaton should be removed by the renormalization procedure in which the cut-off Λ is replaced by a scale for renormalization R , i.e. the characteristic length of the black hole.

At $(1, 1)$ -order, the equation for $\varphi_{(1,1)}$ have no source terms and is not created. The equation for $A_{(1,1)}^i$ is given by

$$\begin{aligned}\partial_\mu \partial^\mu A_{(1,1)}^i &= \partial_0 \left[\dot{\mathbf{A}}_{(1,0)} \cdot (\nabla \times \mathbf{A}_{(0,0)}) \epsilon^{ij}_k \partial_j A_{(0,0)}^k \right] + \partial_a \left[\dot{\mathbf{A}}_{(1,0)} \cdot (\nabla \times \mathbf{A}_{(0,0)}) \epsilon^{bai} \dot{A}_{(0,0)b} \right] \\ &\quad + \partial_0 \left[\dot{\mathbf{A}}_{(0,0)} \cdot (\nabla \times \mathbf{A}_{(1,0)}) \epsilon^{ij}_k \partial_j A_{(0,0)}^k \right] + \partial_a \left[\dot{\mathbf{A}}_{(0,0)} \cdot (\nabla \times \mathbf{A}_{(1,0)}) \epsilon^{bai} \dot{A}_{(0,0)b} \right] \\ &\quad + \partial_0 \left[\dot{\mathbf{A}}_{(0,0)} \cdot (\nabla \times \mathbf{A}_{(0,0)}) \epsilon^{ij}_k \partial_j A_{(1,0)}^k \right] + \partial_a \left[\dot{\mathbf{A}}_{(0,0)} \cdot (\nabla \times \mathbf{A}_{(0,0)}) \epsilon^{bai} \dot{A}_{(1,0)b} \right].\end{aligned}\quad (45)$$

This equation describes four point interaction of photons. Similarly as the case of $(1, 0)$ -order photon (37), one of the $(0, 0)$ -photon $A_{(0,0)}^i$ is chosen to the negative energy photon in the right hand side so that Eq.(45) can be interpreted as two photon scattering. These scattered $(1, 1)$ -order photon $A_{(1,1)}$ and $(0, 0)$ -order photon $A_{(0,0)}^i$ are annihilated to produce $(2, 1)$ -order dilaton as follows,

$$\begin{aligned}(-\partial_t^2 + \partial_r^2 - \frac{15}{4r^2})\varphi_{(2,1)} &= \\ -\delta(r) \frac{\sqrt{2}}{\pi^3 r^{5/2}} \partial_\mu A_{(0,0)i} \partial^\mu A_{(1,1)}^i.\end{aligned}\quad (46)$$

The source term of this equation is integrated over the D3-brane to give

$$\begin{aligned}\frac{1}{V_3} \int d^3x \partial_\mu A_{(0,0)i} \partial^\mu A_{(1,1)}^i &= \\ \frac{C \omega^{10}}{3 \cdot 2^{11} \sqrt{2} \pi^4} e^{-i\omega t} \left(\pi^2 + 2\pi i \log \frac{\omega}{\Lambda} - \left(\log \frac{\omega}{\Lambda} \right)^2 \right).\end{aligned}\quad (47)$$

Using Green's function for a dilaton Eq.(43), we obtain

the solution of Eq.(46) integrated over the D3-brane,

$$\begin{aligned}\bar{\varphi}_{(2,1)} &= \frac{1}{V_3} \int_{V_3} d^3x \varphi_{(2,1)} \\ &= -\int dt' dr' G_R \frac{\sqrt{2} \delta(r')}{\pi^3 r'^{5/2}} \frac{1}{V_3} \int d^3x \partial_\mu A_{(0,0)i} \partial^\mu A_{(1,1)}^i \\ &= C \frac{\omega^{12} e^{-i\omega t}}{3 \cdot 2^{14} \pi^4} H_2^{(1)}(\omega r) \sqrt{r} \\ &\quad \times \left(i - \frac{2}{\pi} \log \frac{\omega}{\Lambda} - \frac{i}{\pi^2} \left(\log \frac{\omega}{\Lambda} \right)^2 \right),\end{aligned}\quad (48)$$

where we again introduce a symmetric factor $1/4$, since the double counting in the momentum integration, which emerged in the evaluation of Eq.(44), occurs twofold in this case. The cut-off Λ will be replaced by the inverse of the characteristic length of the black hole R^{-1} by the procedure of renormalization. The obtained wave function $\bar{\varphi}_{(2,1)}$ correspond to the external line corrected by the 2-loop Feynman diagram depicted in Fig.2.

Assembly of the calculated dilaton waves at several




FIG. 2: The 2-loop self energy for the dilaton. A straight and wavy lines represent propagating dilaton and photon, respectively

orders leads to

$$\bar{\varphi} = \bar{\varphi}_{(0,0)} + \kappa_{10}^2 \bar{\varphi}_{(2,0)} + \frac{\kappa_{10}^2}{T_3} \bar{\varphi}_{(2,1)} + \dots \quad (49)$$

$$= C e^{-i\omega t} \sqrt{r} \left[\mathcal{R} H_2^{(1)}(\omega r) + H_2^{(2)}(\omega r) \right] \quad (50)$$

$$\rightarrow C e^{-i\omega t} \sqrt{\frac{2}{\pi\omega}} \left[\mathcal{R} e^{i(\omega r - \frac{5}{4}\pi)} + e^{-i(\omega r - \frac{5}{4}\pi)} \right], \quad (51)$$

where in the last line, we take $r \rightarrow \infty$ limit and $\mathcal{R}$ is the reflective amplitude,

$$\begin{aligned} \mathcal{R} &= 1 - \frac{\kappa_{10}^2 \omega^8}{2^{11} \pi^3} \left(1 + \frac{i}{\pi} \log \omega R \right) \\ &+ \frac{\kappa_{10}^2}{T_3} \frac{\omega^{12}}{3 \cdot 2^{18} \pi^4} \left(i - \frac{2}{\pi} \log \omega R - \frac{i}{\pi^2} (\log \omega R)^2 \right) + \dots \end{aligned} \quad (52)$$

Here, we recall the relation between κ_{10} , T_3 and R

$$\kappa_{10} = \frac{\sqrt{\pi}}{T_3}, \quad R^4 = \frac{\kappa_{10}}{2\pi^{5/2}}, \quad (53)$$

where we set $N = 1$. Thus the condition of QNM discussed in the appendix A is given by

$$\begin{aligned} 0 &= \frac{\mathcal{I}}{\mathcal{R}} e^{i\beta'} \\ &= 1 + \frac{\pi^2}{512} (\omega R)^8 \left(1 + \frac{i}{\pi} \log \omega R \right) \\ &+ \frac{\pi^2}{512 \cdot 12} (\omega R)^{12} \left(\log \omega R + \frac{i}{2\pi} (\log \omega R)^2 \right) \\ &+ \mathcal{O}((\omega R)^{16}), \end{aligned} \quad (54)$$

where β' is a phase factor,

$$\beta' = \frac{\pi^3}{512 \cdot 24} (\omega R)^{12}. \quad (55)$$

Eq.(54) up to $(\omega R)^8$ is very similar with Eq.(15), and the solution has the qualitatively equivalent behavior.

IV. DISCUSSION

We have evaluated quasi-normal modes (QNMs) of the D3-brane black hole by two different approach, the one is based on the black hole picture in terms of the black

3-brane solution of the low-energy supergravity action and the other the D3-brane described by the field theory with the Dirac-Born-Infeld action. We have shown that the two different method derived qualitatively same condition for the QNMs for the low-energy region. As described in appendix C, the evaluations of absorption cross section shows similar agreement. In both cases, difference, which appears in logarithmic terms, seems to imply that we must take proper account of the non-abelian nature of the theory[11, 12, 34].

The condition for QNMs (A10) has been obtained as Eq.(13) and Eq.(54) in terms of an expansion with respect to ωR . Up to the order $(\omega R)^8$, we can express the condition of QNM as follows,

$$\frac{\mathcal{I}}{\mathcal{R}} \sim 1 + i\Sigma = 0, \quad (56)$$

where $\sim$ means equality up to phase factors and

$$\Sigma = \frac{\pi^2}{512} (\omega R)^8 \left(\frac{\alpha}{\pi} \log \frac{\omega \bar{R}}{2} - i \right). \quad (57)$$

In the field theoretical approach as in section 3, Σ , which is just the coefficient of $\phi_{(2,0)}$, corresponds to the 1-loop self energy of dilaton. The procedure of the spacetime approach gives the dilaton wave function as (51), and we can read off reflective amplitude (52) which is expressed as

$$\frac{\mathcal{R}}{\mathcal{I}} \sim 1 - i\Sigma \quad (58)$$

up to the order $(\omega R)^8$. In our calculation this reflective amplitude is perturbatively inverted to give Eq.(56). However, in the viewpoint of evaluation based on field theories, the above inversion can be interpreted as the summation of the self-energy diagrams

$$\frac{\mathcal{R}}{\mathcal{I}} \sim 1 + (-i\Sigma) + (-i\Sigma)^2 + \dots \sim \frac{1}{1 + i\Sigma}. \quad (59)$$

Thus QNMs can be understood as poles of the scattered propagator in the field theory on D3-brane. This is realized in the calculation of AdS/CFT correspondence in the BTZ black hole case in [29]

Acknowledgments

We would like to thank T.Uematsu and S.Yamaguchi for useful discussions.

APPENDIX A: A BRIEF REVIEW OF QNM

In this appendix, we give a brief review of Quasi-normal modes (QNMs) which represent the relaxation process in a black hole spacetime. If one perturbs the black hole spacetime, then gravitational wave is emitted

to both directions. One part of the wave propagates to spatial infinity and other falls into black hole horizon. So, it is natural to specify the boundary condition for such waves as follows, ingoing at horizon and outgoing at spatial infinity.

Let us consider a minimal scalar on the black holes background. It satisfies the wave equation of following forms after some change of variables[15],

$$(-\partial_t^2 + \partial_{r_*}^2 - V(r_*))\phi = 0 \quad (\text{A1})$$

where r_* is the ‘‘tortoise’’ radial coordinate. In the case of the Schwarzschild black hole with the horizon radius $r = r_h$, the coordinate r_* is related to the usual radial coordinate r as

$$r_* = r + r_h \ln\left(\frac{r - r_h}{r_h}\right),$$

which spans the region outside of the horizon of the black hole. It follows that $r_* \rightarrow -\infty$ represents approaching the horizon ($r = r_h$) and $r_* \rightarrow \infty$ the spatial infinity ($r \rightarrow \infty$), respectively. And the potential $V(r_*)$, which has the information of the curvature of the spacetime, is assumed to have the asymptotic behavior, $V(r_*) \rightarrow 0$ as $r_* \rightarrow -\infty$ (to horizon) and $r_* \rightarrow \infty$ (to spatial infinity). Under this assumption, the solution of wave equation (A1) with a frequency ω can be expressed as

$$\phi(t, r_*) = Ae^{-i\omega(t-r_*)} + Be^{-i\omega(t+r_*)} \quad (\text{A2})$$

in the asymptotic region. Here let me remind you the boundary conditions, i.e. ‘‘QNM’s being ingoing at the horizon and outgoing at the spatial infinity’’ are specified by

$$\phi \propto \begin{cases} e^{-i\omega(t-r_*)}, & (r_* \rightarrow \infty) \\ e^{-i\omega(t+r_*)}, & (r_* \rightarrow -\infty) \end{cases} \quad (\text{A3})$$

In order to obtain the solution which satisfies the boundary condition described above, we consider the retarded Green’s function constructed by following prescription. We prepare two solutions $\phi_1(t, r_*)$ and $\phi_2(t, r_*)$, which has asymptotic forms as

$$\phi_1 \sim \begin{cases} \mathcal{I}e^{-i\omega(t+r_*)} + \mathcal{R}e^{-i\omega(t-r_*)}, & (r_* \rightarrow \infty) \\ \mathcal{A}e^{-i\omega(t+r_*)}, & (r_* \rightarrow -\infty) \end{cases} \quad (\text{A4})$$

$$\phi_2 \sim \begin{cases} \mathcal{Q}e^{-i\omega(t-r_*)}, & (r_* \rightarrow \infty) \\ \mathcal{P}e^{-i\omega(t-r_*)} + \mathcal{G}e^{-i\omega(t+r_*)}, & (r_* \rightarrow -\infty) \end{cases} \quad (\text{A5})$$

to construct the retarded Green’s function from them,

$$G_R(t, r_*; t', r'_*) = \theta(t - t') \int \frac{d\omega}{2\pi} e^{-i\omega(t-t')} \frac{1}{\Delta} \times [\theta(r_* - r'_*)\phi_1(r'_*)\phi_2(r_*) + (r_* \leftrightarrow r'_*)], \quad (\text{A6})$$

where the Wronskian Δ can be evaluated as $\Delta = 2i\omega\mathcal{I}\mathcal{Q}$ from the asymptotic forms at $r_* \rightarrow \infty$. Using this retarded Green’s function, we can obtain the wave ϕ generated by a source term $S(t, r_*)$

$$\phi(t, r_*) = \int dt' dr'_* G_R(t, r_*; t', r'_*) S(t', r'_*), \quad (\text{A7})$$

which satisfies the inhomogeneous equation,

$$(-\partial_t^2 + \partial_{r_*}^2 - V(r_*))\phi = S(t, r_*). \quad (\text{A8})$$

It is obvious that the solution (A7) satisfies the QNM boundary condition (A3). Taking the limit $r_* \rightarrow \infty$ of Eqs. (A7) and (A6), we obtain the asymptotic form of the generated wave

$$\phi(t, r_*) \rightarrow \int_C dt' dr'_* \theta(t - t') \int_C \frac{d\omega}{2\pi} \times \frac{e^{-i\omega(t-t')} e^{i\omega r_*}}{2i\omega\mathcal{I}} \phi_1(r'_*) S(t', r'_*), \quad (\text{A9})$$

where contour C must be taken as it enclose lower half of complex ω plane for retarded boundary condition. So, only the poles of lower half plane contribute the ω integral in the asymptotic form Eq.(A9) and that is the Quasi-normal modes (QNM) which are, of course, complex and decaying modes. In other words, QNM is the poles of retarded Green’s function on the lower half complex ω plane. The condition of QNM is $\mathcal{I} = 0$, but only the ratio of amplitudes are meaningful in the asymptotic form Eq.(A4), so this is equivalent to $\mathcal{R} \rightarrow \infty$. Then the more general condition of QNM is,

$$\mathcal{I}/\mathcal{R} = 0 \quad (\text{A10})$$

The only things we have to do is to solve this equation and choose modes arising at lower half complex plane.

APPENDIX B: EXPLICIT EXPRESSIONS

In this appendix, we summarize explicit form for some results which were considered too lengthy to write out in the main text.

The Floquet exponent μ which satisfies Eq.(8) can be obtained in the expansion with respect to a power series in ωR ,

$$\mu = 1 - \frac{i\sqrt{5}}{6}(\frac{\omega R}{2})^4 + \frac{7i}{216\sqrt{5}}(\frac{\omega R}{2})^8 + \frac{11851i}{62208\sqrt{5}}(\frac{\omega R}{2})^{12} + \dots \quad (\text{B1})$$

And the coefficient $A_\mu^{(q)}$ in Eq.(7) up to $(\omega R)^{16}$ are given by

$$\begin{aligned} A_\mu^{(1)} &= \frac{1}{3\mu(\mu+1)(\mu+2)}, \\ A_\mu^{(2)} &= \frac{1}{432} \left(\frac{1}{\mu} - \frac{11}{\mu+1} - \frac{47}{\mu+2} \right) - \frac{1}{144} \frac{1}{\mu+3} - \frac{1}{18} \frac{1}{(\mu+2)^2} + \frac{5}{36} \psi^{(1)}(\mu+2), \end{aligned} \quad (\text{B2})$$

$$\begin{aligned} A_\mu^{(3)} &= -\frac{2321}{155520\mu} + \frac{739}{9720(\mu+1)} - \frac{5}{216(\mu+2)^3} - \frac{115}{2592(\mu+2)^2} \\ &\quad - \frac{791}{9720(\mu+2)} + \frac{1}{432(\mu+3)^2} + \frac{61}{6480(\mu+3)} + \frac{1}{17280(\mu+4)} \\ &\quad + \left(\frac{7}{648} + \frac{5}{216\mu} - \frac{5}{108(\mu+1)} + \frac{5}{216(\mu+2)} \right) \psi^{(1)}(\mu+2) \\ A_\mu^{(4)} &= \left(\frac{164327}{133996800} - \frac{37\pi^2}{31104} \right) \frac{1}{\mu+1} + \dots, \end{aligned} \quad (\text{B3})$$

where, $\psi^{(1)}$ is the PolyGamma function.

APPENDIX C: ABSORPTION CROSS SECTION FOR D3-BRANE

In this appendix, we compare absorption cross section σ for the D3-brane in various methods for evaluation. As showed in [11], there is a bit difference in σ for different methods at higher order.

In supergravity, the absorption cross section σ for the 3-brane is given by

$$\begin{aligned} \sigma_{\text{sugra}} &= \frac{32\pi^2}{\omega^5} \frac{|\mathcal{A}|^2}{|\mathcal{I}|^2} \\ &= \frac{\pi^4}{8} \omega^3 R^8 \left(1 - \frac{1}{6} (\omega R)^4 \log \omega \bar{R} + \mathcal{O}((\omega R)^4) \right) \end{aligned} \quad (\text{C1})$$

from Eq.(11) as in [12]. This evaluation is reliable at $N \gg 1$ region where the supergravity description of the

3-brane is valid.

We can also calculate the absorption cross section σ for the D3-brane by means of the space-time approach from Eq.(51),

$$\begin{aligned} \sigma_{\text{wave}} &= \frac{32\pi^2}{\omega^5} \left(1 - \left| \frac{\mathcal{R}}{\mathcal{I}} \right|^2 \right) \\ &= \frac{\pi^4}{8} \omega^3 R^8 \left(1 + \frac{1}{12} (\omega R)^4 \log \frac{\omega}{\Lambda} + \mathcal{O}((\omega R)^4) \right), \end{aligned} \quad (\text{C2})$$

where we set $N = 1$.

As in [11], CFT on the D3-brane can also derive this quantity, and result is

$$\sigma_{\text{CFT}} = \frac{\pi^4}{8} \omega^3 R^8 \left(1 - \frac{1}{24} (\omega R)^4 \log \frac{\omega}{\Lambda} + \mathcal{O}((\omega R)^4) \right). \quad (\text{C3})$$

These three evaluation of σ are agree at the lowest order.

-
- [1] J.D.Bekenstein, Phys.Rev.D **5** 1239, (1972)
 - [2] S.W.Hawking Phys.Rev.Lett. **26**, 1344 (1971)
 - [3] S.W.Hawking Commun.Math.Phys. **43** 199, (1975)
 - [4] e.g. R.M.Wald “*General Relativity*” (The University of Chicago Press, Chicago and London, 1984)

- [5] e.g. N.D.Birrell and P.C.W.Davies “*Quantum Fields in Curved Space*” (Cambridge University Press, Cambridge, 1982)
- [6] See e.g. G.Horowitz gr-qc/9604051.
- [7] A.Strominger and C.Vafa Phys. Lett. B **379**, 99 (1996).

- [8] C.G.Callan and J.M.Maldacena Nucl.Phys.**B472**,591 (1996).
- [9] S.R.Das and S.D.Mathur Nucl.Phys.**B 478**, 561 (1996).
- [10] I. R. Klebanov Nucl. Phys. **B 496**, 231 (1997).
- [11] S. S. Gubser, A. Hashimoto, I. R. Klebanov and M. Krasnitz Nucl. Phys. **B 526** , 393 (1998).
- [12] S. S. Gubser and A. Hashimoto hep-th/9805140
- [13] Proc. R. Soc. London, **A 344** 441 (1975).
- [14] See e.g. S. Chandrasekhar, *The Mathematical Theory of Black Holes* (Clarendon, Oxfrd, 1983)
- [15] See e.g. K.D.Kokkotas and B.G. Schmidt Living Reviews in Relativity "Quasi-normal modes of Stars and Black Holes", (<http://www.livingreviews.org/Articles/Volume2/1999-2kokkotas/index.html>)
- [16] G.T.Horowitz and V.E.Hubeny Phys.Rev.D **62** (2000) 024027
- [17] J.S.F. Chan, R.B. Mann Phys.Rev.D **55** (1997) 7546
- [18] J.S.F. Chan, R.B. Mann Phys.Rev.D **59** (1999) 064025
- [19] B.Wang, C.Lin, E.Abdalla Phys.Lett.B **481** (2000) 79-88
- [20] T.R.Govindarajan, V.Suneeta Class.Quant.Grav.**18** (2001) 265-276
- [21] Jiong-Ming Zhu, Bin Wang, Elcio Abdalla Phys.Rev. D **63** (2001) 124004
- [22] Danny Birmingham Phys.Rev. D **64** (2001) 064024
- [23] Vitor Cardoso, Jose' P. S. Lemos Phys.Rev. D **64** (2001) 084017
- [24] Vitor Cardoso, Jose' P. S. Lemos Class.Quant.Grav.**18** (2001) 5257-5267
- [25] Ian G Moss, James P Norman Class.Quant.Grav. **19** (2002) 2323-2332
- [26] B. Wang, E. Abdalla, R.B. Mann Phys.Rev.D **65** (2002) 084006
- [27] R.A.Konoplya hep-th/0205142, gr-qc/0207028
- [28] Andrei O. Starinets hep-th/0207133
- [29] D.Birmingham, I.Sachs and S.N.Solodukhin Phys.Rev.Lett **88**, 151301 (2002).
- [30] R. P. Feynman Rev. Mod. Phys. **21**, 425 (1949).
- [31] R. P. Feynman Rev. Mod. Phys. **20**, 367 (1948).
- [32] R. P. Feynman Phys. Rev. **76**, 749 (1949).
- [33] J. J. Sakurai "Advanced Quantum Mechanics" Addison Wesley Publishing Company **Chapter 4-5**
- [34] C.G.Callan,S.S.Gubser,I.R.Klebanov,and A.A.Tseytlin, Nucl.Phys. **B 489**,65 (1997).